

This paper is “in press” (Perception)

Judging body-weight from faces: The height-weight illusion

Tobias M. Schneider¹,

Heiko Hecht¹ & Claus-Christian Carbon^{2,*}

¹Department of Psychology, Johannes Gutenberg-University Mainz

²Department of General Psychology and Methodology, University of Bamberg

*) **Corresponding author:** ccc@experimental-psychology.com

Keywords: face-specific processing; weight illusion; height; viewing angle; distortion perception bias; vantage point; perspective; health; indicator

Abstract

Being able to exploit features of the human face to predict health and fitness can serve as an evolutionary advantage. Surface features such as facial symmetry, averageness, and skin colour are known to influence attractiveness. We sought to determine whether observers are able to extract more complex features, namely body weight. If possible, it could be used as a predictor for health and fitness. For instance, facial adiposity could be taken to indicate a cardiovascular challenge or proneness to infections. Observers seem to be able to glean body weight information from frontal views of a face. Is weight estimation robust across different viewing angles? We showed that participants strongly overestimated body weight for faces photographed from a lower vantage point while underestimating it for faces photographed from a higher vantage point. The perspective distortions of simple facial measures (e.g. width-to-height ratio) that accompany changes in vantage point do not suffice to predict body weight. Instead, more complex patterns must be involved in the height-weight illusion.

Judging body-weight from faces: The height-weight illusion

Besides variables such as facial symmetry (e.g. Grammer and Thornhill 1994), averageness (e.g. Carbon et al 2010) and skin colour (e.g. Jones et al 2004), perceived human weight proves to be a reliable predictor for health and fitness (Coetzee et al 2009; Swami et al 2008). Perceived body weight also plays a significant role in judgments of attractiveness (Thornhill and Grammer 1999; Tovee et al 1998). Importantly, people make fairly accurate weight judgments on the basis of standardized facial images.

We do not often view faces from a vantage point centred on and aligned with the face. Are weight estimations still reliable and valid with changing vantage points? We investigated the impact of viewing angle on perceived human body weight in 2D pictures. In order to ascertain a precise orientation of the face with respect to the vantage point of the camera, we selected 3D face scans (di3d-technology) of 48 human models (24 female, aged $M=23.5$ ys, $SD=16.0$, range 3 to 56 ys), aligned them with respect to a virtual camera, and created 2D images of the faces corresponding to a camera position aligned with the inter-ocular point and perpendicular to the vertical axis of the face. We then rendered the image from three camera heights (see Figure 1): -30° (elevated, which is equivalent to the head being pitched downwards by -30° or a raised and tilted camera), 0° (frontal view), and $+30^\circ$ (lowered, which is equivalent to the head pitched upwards by $+30^\circ$ or a lowered and tilted camera). We refer to this manipulation as *viewing angle* in the following.

insert Figure 1 about here

Participants (18 female, $M=36.8$ ys, $SD=17.2$, range 18 to 69 ys) were exposed to all 48 faces. The experiment had two main factors which were hierarchically organized with *model gender* (gender of the shown face) as the superordinate and *viewing angle* as the

subordinate order. Factors' levels were blocked and the sequences of them were counterbalanced across participants. The observers' task was to judge the body weight of the faces shown in kg.

Weight estimations showed very high reliabilities across participants (Table 1), which matches previous findings (Coetzee et al 2009). A two-way by-items mixed-design ANCOVA with the between-subjects variable *model gender*, the within-subjects variable *viewing angle* (-30°, frontal, +30°) and the covariate *model age* (age of the face models) was conducted. *Viewing angle* showed a large effect, $F(2,90)=78.1$, $p<.0001$, $\eta_p^2=.634$, revealing a strong influence of changes in vantage point: The higher the observer, the lighter the judged body weight. We also found effects of *model gender*, $F(1,45)=5.13$, $p=.0284$, $\eta_p^2=.102$, and *model age*, $F(1,45)=144.04$, $p<.0001$, $\eta_p^2=.762$, which was not surprising owing to the fact that female and young faces are commonly associated with lower body weight.

insert Table 1 about here

How can this height-weight illusion be explained? Proportion changes in the projected image could serve as a cue to body weight, such as the width-ratio of upper versus lower face. A lower vantage point causes a relatively wider lower face in the projected image. To test if these geometric changes are associated with greater body-weight, we measured cheek-to-jaw-width ratio, width-to-upper-height ratio and width-to-lower-height ratio following Coetzee et al's (2010) suggestion (see Figure 2 and Table 2 for results).

insert Figure 2 about here

insert Table 2 about here

Only the correlation between the ratio of width-to-upper-facial-height and weight was significant for the *viewing angles* up vs. frontal ($r=.415, p<.01$), all other correlations between relative geometric changes and weight estimation were not. However, the effect of *viewing angle* is not tightly predicted by the change of these geometrical cues. Observers appeared to be able to partially compensate for the changes in perspective geometry. For instance, when lowering the camera the average weight change (+ 22 %) was associated with an even larger change in width-to-upper-facial-height ratio (+ 30 %). Thus, other factors than straightforward projective geometry must be involved in the height-weight illusion.

Figures

Figure 1. Illustration of used *viewing angles* (e.g. $+30^\circ$: observer's vantage point is below the observed face).





Figure 2. Measures used to calculate facial cues according to Coetzee et al (2010). (a) Width-to-lower-facial-height ratio: Cheekbone width divided by lower facial height. (b) Width-to-upper-facial-height ratio: Cheekbone width divided by upper facial height. (c) Cheek-to-jaw-width ratio: Cheekbone width divided by jaw width.



Tables

Table 1. Average weight estimations and reliabilities (Cronbach's α) for different *viewing angles* split by *model gender*.

Viewing angle	<i>M</i>	<i>SD</i>	<i>Range</i>	Cronbach's α
Female				
-30°	41.10	4.42	[24.88-47.98]	.986
Frontal	50.22	3.61	[38.88-57.56]	.993
+30°	61.07	5.07	[52.50-77.50]	.990
Male				
-30°	49.78	4.94	[27.75-55.16]	.992
Frontal	58.96	3.81	[50.13-66.25]	.996
+30°	72.26	5.02	[62.13-81.94]	.988

Table 2. Parameters of facial geometry (averaged across all faces) by *viewing angle*.

Distances are measured in pixels.

Viewing angle	<i>M</i>	<i>SD</i>	<i>M</i>	<i>SD</i>	<i>Range</i>
	Cheekbone width		Cheekbone-to-jaw-width ratio		
-30°	274.35	23.42	1.62	0.13	[1.35-1.88]
Frontal	324.17	25.25	1.73	0.05	[1.10-1.32]
+30°	274.73	21.88	0.95	0.03	[0.87-1.02]
	Upper facial height		Width-to-upper-height ratio		
-30°	139.69	15.88	1.97	0.12	[1.72-2.30]
Frontal	168.71	25.68	1.95	0.19	[1.58-2.33]
+30°	111.54	21.54	2.53	0.35	[1.95-3.46]
	Lower facial height		Width-to-lower-height ratio		
-30°	65.02	8.97	4.27	0.50	[3.44-5.80]
Frontal	124.01	14.95	2.63	0.22	[2.20-3.10]
+30°	138.19	17.17	2.01	0.18	[1.55-2.38]
	Jaw width				
-30°	170.25	17.98			
Frontal	276.67	22.21			
+30°	289.83	23.71			

References

- Carbon C C, Gruter T, Gruter M, Weber J E, Lueschow A, 2010 "Dissociation of facial attractiveness and distinctiveness processing in congenital prosopagnosia" *Visual Cognition* **18** 641-654
- Coetzee V, Chen J Y, Perrett D I, Stephen I D, 2010 "Deciphering faces: Quantifiable visual cues to weight" *Perception* **39** 51-61
- Coetzee V, Perrett D I, Stephen I D, 2009 "Facial adiposity: A cue to health?" *Perception* **38** 1700-1711
- Grammer K, Thornhill R, 1994 "Human (Homo sapiens) facial attractiveness and sexual selection: The role of symmetry and averageness" *Journal of Comparative Psychology* **108** 233-242
- Jones B C, Little A C, Burt D M, Perrett D I, 2004 "When facial attractiveness is only skin deep" *Perception* **33** 569-576
- Swami V, Miller R, Furnham A, Penke L, Tovee M J, 2008 "The influence of men's sexual strategies on perceptions of women's bodily attractiveness, health and fertility" *Personality and Individual Differences* **44** 98-107
- Thornhill R, Grammer K, 1999 "The body and face of woman: One ornament that signals quality?" *Evolution and Human Behavior* **20** 105-120
- Tovee M J, Reinhardt S, Emery J L, Cornelissen P L, 1998 "Optimum body-mass index and maximum sexual attractiveness" *Lancet* **352** 548-548

Author's note

We would like to thank Vera M. Hesslinger and Sabine Albrecht at the Department of General Psychology and Methodology, University of Bamberg for assisting in data analysis.